



EDWARD R. ANDERSON, MD
1830 N. Franklin St, Suite 340
Denver, CO 80218
PHONE: (303) 318-3330 FAX: (303) 812-4221

www.FROST-clinic.com

**COMPREHENSIVE ADULT SPINE, ADULT TOTAL HIP AND KNEE RECONSTRUCTION, AND
ORTHOPAEDIC TRAUMATOLOGY/FRACTURE MANAGEMENT**

Patient Name: _____

Diagnosis: _____

Lumbar Disc Replacement Physical Therapy Prescription

The intent of this protocol is to provide guidelines for rehab. It is not intended to be used as a substitute for clinical decision-making.

If any of the following occur, contact Dr. Anderson and hold off on physical therapy:

- Any signs of infection
- Worsening of radicular symptoms, including progressive weakness
- Unexpected high self-reports of pain in comparison to presurgical state

Phase I (0-2 Weeks): Protective Phase

Therapy

- First visit at 2 weeks postoperatively as an outpatient.

Precautions

- Avoid bending, lifting, and twisting (**No BLT**). No pushing and pulling. Nothing heavier than 5lbs until 6 weeks postoperatively.
- Limit sitting, including in the car, to no more than 30 minutes at a time (standing/walking breaks - at least 50-100 feet at a time).
- No extension range of motion, and no rotation exercises for 8 weeks postoperatively.

Goals:

- Diminish pain/inflammation and minimize lower extremity radiating symptoms (ice, modalities as needed).
- Learn correct posture, body mechanics, transfers, and positioning.
- Achieve proper muscle firing for transversus abdominis, multifidus, and gluteal motor groups
- Focus on walking program, increasing tolerance to 10 minutes, 2 times a day.

Education

- **Posture Education:** Upright sitting posture with Lumbar Roll at all times; Frequent changes in positions, and sleeping positions. Teach how different body positions and postures affect the spine.
- **Body Mechanics:** Light lifting, transfers (include logrolling), positioning, etc.

Exercises

- **Walking Program:** Begin 1-2 times a day for 10 minutes, progress as tolerated
- **Transversus Abdominis Bracing:** 10 second isometrics with normal breathing (without pelvic tilt) in supine position.
- **Multifidus:** 10 second isometrics with normal breathing in prone position (if able).
- **Glute sets:** 10 second isometrics with emphasis on proper gluteal muscle group firing in prone position (bilateral and unilateral).
- **Light stretching:** Hip flexors, quadriceps, hamstrings, and gastrocsoleus.

- **Driving:** When off narcotic pain medications, and patient feels that they are safe enough to drive, able to react to avoid accidents/drive defensively without hesitation. Dr. Anderson does not clear patients to drive, this is an individual decision for each patient.

Phase II (2-6 Weeks): Initial Strengthening Phase

Therapy

- Starting at week 2, 1-2 times per week, 4 or more weeks

Precautions

- Keep spine in neutral and good posture for strengthening with focus on proper neuromuscular control, do not progress without good neuromuscular control.
- **Lifting Restrictions:** Lifting, pushing and pulling less than **5-10 pounds** until 6 weeks.
- No extension range of motion, and no rotation exercises for 8 weeks.

Goals

- Patient to have proper neuromuscular control and posture with stabilization and strength exercises
- Complete light strengthening, with a neutral spine and correct firing of stabilizers.
- Release soft tissue restriction/muscle spasm/scar
- Independent with body and lifting mechanics - review
- Increase aerobic endurance to more than 30 minutes

Cardiovascular (as long as it does not aggravate or increase symptoms, let pain be your guide)

- Walking Progression: Gradually progress to 30 minutes a day.
- Biking – Recumbent Stationary Bike (avoid road bikes) initiate at 2 weeks postop
- Stationary Bike Upright - Can initiate at 4 weeks (no resistance), 6 weeks (resistance)

Strength

Only initiate Phase II strength conditioning once patient can complete Phase I exercises. Then begin light resistance and slowly progress. Emphasize achieving proper neuromuscular control of the transversus abdominis and multifidus muscle groups without compensation (use stabilizer biofeedback cuff if available).

- **Transversus Abdominis/Multifidus Progression:** 10-20 second isometrics with normal breathing (without pelvic tilt) in supine, quadruped and prone. Must maintain neutral spine.
 - Progress with upper extremity/lower extremity movements (example marches, straight leg raises, upper extremity lift and lowers, planks, etc.)

- **Supine:** Add upper extremity/lower extremity movements (example: Marches, straight leg raises, upper extremity lift and lowers, etc.)
- **Quadruped:** Alternating upper extremity, alternating lower extremity, alternating upper extremity/lower extremity (bird-dog)
- **Bridging:** Bridges, Side Leg Bridges, on Swiss ball, etc.
- **Balance:** Static (example: Single-Leg Stance, Tandem, etc.)
- **Swiss ball:** No Sit Ups, No Twisting (example wall squats, seated exercises, bridges, bird-dog, etc.)
- **Multifidus:** 10-20 second isometrics with normal breathing in prone position.
- **Glute Activation Exercises:** 10-20 second isometrics with emphasis on proper gluteal muscle group firing in prone position (bilateral and unilateral).
 - **Prone:** Hip extensions, alternating upper extremity/lower extremity lifts (avoid hyperextension)
 - **Side-lying:** Clams, hip abduction, etc
 - 90/90 leg lifts, side-lying abduction, quadruped hip extension, bird-dog
 - **Bridging:** Bridges, Side Leg Bridges, on Swiss ball, etc.
- **Upper Extremity/Lower Extremity Strength Training**
 - **Step ups, leg press, wall squats, body weight squats, etc.**
 - **Balance:** With transversus abdominis bracing): Single-leg stance, tandem, foam, etc.
 - **Upper extremity light resistive exercises** (machines, Thera-Band, free weights).

Flexibility

- **Stretching:** Hamstrings, gastroc/soleus, quadriceps, hip flexors, piriformis, etc.
- **Neural mobilization:** Performed as needed, gentle with caution not to flareup nerve roots

Aquatic Physical Therapy (At 4 weeks if available once incision is healed)

- No rotation and transversus abdominis bracing during all exercises
- Walking all directions, balance, upper extremity/lower extremity strengthening

Phase III (6-8 weeks): Progression to Advanced Strengthening

Therapy

- 1-2 times a week (as needed for return to sport or work)

Precautions

- No extension range of motion, and no rotation exercises for 8 weeks postoperatively.

Goals

- Independent home exercise program for advance strengthening, return to sport and work.
- Increased lower quarter flexibility and strength with focus on proper transversus abdominis and gluteal activation.
- Anticipate release to full activities without restrictions at the 6-8 week post operative time point. Pending approval by Dr. Anderson.
- Possible referral to work hardening/reconditioning program.

Strength

- Advanced Core Strength and Stabilization Exercises:
 - **Highly Recommend Pilates**
 - Progress to weightbearing, balance, Swiss Ball, reformer, etc.

- Progressed to multiplanar exercises with upper and lower extremities
- Progress lower extremity/upper extremity strengthening
- Begin running, agility and plyometric exercise program for return to sport at 8 to 12 weeks (if symptoms are stable and patient is cleared by Dr. Anderson)
- Possible referral to Work Hardening/Work Reconditioning program

Flexibility

- Lumbar spine more than 8 weeks: Work on improving lumbar extension range of motion, but avoid end range (example: Prone lying, prone on elbows, press ups and/or gentle standing exercises if no peripheralization)

Cardiovascular

- As long as it does not aggravate or increase symptoms, let pain be your guide
- Walking - The best exercise for these patients!
- Biking - Recumbent or hybrid/upright (avoid road bikes)
- Swimming
- Elliptical
- Hiking
- Pilates
- Emphasize correct form and equipment set up (example: Elliptical, bike, walking training, etc.)
- Preference of Pilates or yoga, if returning to yoga, and sure it is with an experienced instructor. Same goes for Pilates.
- When initiating running and sports below, slowly increase in the 8 to 12-week timeframe.
- Time frames may vary per patient, and based upon the sport/activity, consult with Dr. Anderson if you have questions.

No Earlier Than:

Walking Progression	At least 30 mins/day, continue progress
Stationary Bike	Gradual increase in resistance at 4 weeks
Pilates (neutral spine)	6 weeks
Outdoor Biking	6 weeks
Hiking	6 weeks
Elliptical	6 weeks
Skiing	8 weeks

No Earlier Than:

Yoga	8 weeks
Swimming	6-8-week progression
Running	8-12 wk progression
Soccer	8-12 wk progression
Basketball	8-12 wk progression
Golf	8-12 wk progression